

Recommended Analgesic Regimens for MICE by Procedure Severity

| Mild Pain | | (e.g., tail biopsy of mice >21 days of age, needle biopsy, percutaneous pellet implantation) | | |
|-----------|------------|--|-----------|-------------|
| Drug | Dose mg/kg | Route | Frequency | Duration |
| meloxicam | 5 - 10 | SQ, PO | once | 12-24 hours |
| carprofen | 5 - 20 | SQ | once | 12-24 hours |

| Moderate Pain | | Minor surgical procedures (e.g., subcutaneous pump implantation, simple vascular cut down) | | |
|------------------|------------|--|---------------|---------------------|
| Drug | Dose mg/kg | Route | Frequency | Duration |
| meloxicam | 5 - 10 | SQ, PO | Q 12-24 hours | 12-24 hours |
| meloxicam ER | 5 | SQ | Q 24 hours | 1-3 days, as needed |
| carprofen | 5 - 20 | SQ | Q 12 hours | 12-24 hours |
| buprenorphine | 0.05 - 0.2 | SQ | Q6-8 hours | 24 hours |
| buprenorphine SR | 0.5 - 1.0 | SQ | once | 24 hours |
| bupivacaine* | 2 | SQ at incision | once | 4-6 hours |

| Severe Pain | | Major surgical procedures (e.g., laparotomy, thoracotomy, craniotomy, extensive tissue dissection) | | |
|------------------|--------------|--|------------------------------------|---------------------|
| Drug | Dose (mg/kg) | Route | Frequency | Duration |
| meloxicam | 5 - 10 | SQ, PO | Q24 hours | 48 hours |
| meloxicam ER | 5 | SQ | Q 24 hours | 3-5 days, as needed |
| carprofen | 5 - 20 | SQ | Q12 hours, Q24 w/ buprenorphine | 48 hours |
| buprenorphine* | 0.1 - 2 | SQ | once | 3-5 hours |
| buprenorphine SR | 0.5 - 1.0 | SQ | Q. 72 hrs | 3-5 days, as needed |
| bupivacaine* | 2 | SQ at incision | once | 4-6 hours |

* Agents are recommended in addition to the listed NSAIDs as a part of a multimodal analgesic regimen.

Mouse Analgesic Dosages

| | Meloxicam | Meloxicam ER* | Buprenorphine | Buprenorphine SR* | Carprofen | Nocita 13.3 |
|-----------------------------|--------------------------|---------------|---------------|-------------------|------------------------------|-----------------------------------|
| Stock concentration (mg/ml) | 5 | 2 | 0.3 | 1 | 50 | |
| instructions for diluting | 1ml meloxicam + 9ml NaCl | n/a | n/a | n/a | 0.2ml carprofen + 9.8ml NaCl | 1 ml Nocita + 1 ml sterile saline |
| Diluted concentration for | 0.5 | n/a | n/a | n/a | 1 | 6.65 |
| Dosage (mg/kg) | 5 - 10 | 5 | 0.05 - 0.2 | 1 | 5 - 20 | 5.3 |
| Frequency | q 12 hrs | q 24 hrs | q 6-8 hrs | q 72 hrs | q 12 hrs | once |
| MOUSE weight in grams | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) |
| | 0.18 - 0.35 | 0.05 | 0.003 - 0.01 | 0.02 | 0.09 - 0.36 | 0.014 |
| | 0.19 - 0.38 | 0.05 | 0.003 - 0.01 | 0.02 | 0.10 - 0.38 | 0.015 |
| | 0.20 - 0.40 | 0.05 | 0.003 - 0.01 | 0.02 | 0.10 - 0.40 | 0.016 |
| | 0.21 - 0.42 | 0.05 | 0.004 - 0.01 | 0.02 | 0.11 - 0.42 | 0.017 |
| | 0.22 - 0.44 | 0.06 | 0.004 - 0.01 | 0.02 | 0.11 - 0.44 | 0.018 |
| | 0.23 - 0.46 | 0.06 | 0.004 - 0.02 | 0.02 | 0.12 - 0.46 | 0.018 |
| | 0.24 - 0.48 | 0.06 | 0.004 - 0.02 | 0.02 | 0.12 - 0.48 | 0.019 |
| | 0.25 - 0.50 | 0.06 | 0.004 - 0.02 | 0.03 | 0.13 - 0.50 | 0.020 |
| | 0.26 - 0.52 | 0.07 | 0.004 - 0.02 | 0.03 | 0.13 - 0.52 | 0.021 |
| | 0.27 - 0.54 | 0.07 | 0.005 - 0.02 | 0.03 | 0.14 - 0.54 | 0.022 |
| | 0.28 - 0.56 | 0.07 | 0.005 - 0.02 | 0.03 | 0.14 - 0.56 | 0.022 |
| | 0.29 - 0.58 | 0.07 | 0.005 - 0.02 | 0.03 | 0.15 - 0.58 | 0.023 |
| | 0.30 - 0.60 | 0.08 | 0.005 - 0.02 | 0.03 | 0.16 - 0.60 | 0.024 |
| | 0.31 - 0.62 | 0.08 | 0.005 - 0.02 | 0.03 | 0.16 - 0.60 | 0.025 |
| | 0.32 - 0.64 | 0.08 | 0.005 - 0.02 | 0.03 | 0.17 - 0.64 | 0.026 |
| | 0.33 - 0.66 | 0.08 | 0.006 - 0.02 | 0.03 | 0.17 - 0.66 | 0.026 |
| | 0.34 - 0.68 | 0.09 | 0.006 - 0.02 | 0.03 | 0.17 - 0.68 | 0.027 |
| | 0.35 - 0.70 | 0.09 | 0.006 - 0.02 | 0.04 | 0.18 - 0.70 | 0.028 |

*Use 23G needles when using ER/SR analgesics

| | Meloxicam | Meloxicam ER* | Buprenorphine | Buprenorphine SR* | Carprofen | Nocita 13.3 |
|-----------------------------|--------------------------|---------------|---------------|-------------------|------------------------------|-----------------------------------|
| Stock concentration (mg/ml) | 5 | 2 | 0.3 | 1 | 50 | |
| instructions for diluting | 1ml meloxicam + 9ml NaCl | n/a | n/a | n/a | 0.2ml carprofen + 9.8ml NaCl | 1 ml Nocita + 1 ml sterile saline |
| Diluted concentration for | 0.5 | n/a | n/a | n/a | 1 | 6.65 |
| Dosage (mg/kg) | 5 - 10 | 5 | 0.05 - 0.2 | 1 | 5 - 20 | 5.3 |
| Frequency | q 12 hrs | q 24 hrs | q 6-8 hrs | q 72 hrs | q 12 hrs | once |
| MOUSE weight in grams | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) |
| | 18 | 0.18 - 0.35 | 0.05 | 0.003 - 0.01 | 0.02 | 0.09 - 0.36 |
| | 19 | 0.19 - 0.38 | 0.05 | 0.003 - 0.01 | 0.02 | 0.10 - 0.38 |
| | 20 | 0.20 - 0.40 | 0.05 | 0.003 - 0.01 | 0.02 | 0.10 - 0.40 |
| | 21 | 0.21 - 0.42 | 0.05 | 0.004 - 0.01 | 0.02 | 0.11 - 0.42 |
| | 22 | 0.22 - 0.44 | 0.06 | 0.004 - 0.01 | 0.02 | 0.11 - 0.44 |
| | 23 | 0.23 - 0.46 | 0.06 | 0.004 - 0.02 | 0.02 | 0.12 - 0.46 |
| | 24 | 0.24 - 0.48 | 0.06 | 0.004 - 0.02 | 0.02 | 0.12 - 0.48 |
| | 25 | 0.25 - 0.50 | 0.06 | 0.004 - 0.02 | 0.03 | 0.13 - 0.50 |
| | 26 | 0.26 - 0.52 | 0.07 | 0.004 - 0.02 | 0.03 | 0.13 - 0.52 |
| | 27 | 0.27 - 0.54 | 0.07 | 0.005 - 0.02 | 0.03 | 0.14 - 0.54 |
| | 28 | 0.28 - 0.56 | 0.07 | 0.005 - 0.02 | 0.03 | 0.14 - 0.56 |
| | 29 | 0.29 - 0.58 | 0.07 | 0.005 - 0.02 | 0.03 | 0.15 - 0.58 |
| | 30 | 0.30 - 0.60 | 0.08 | 0.005 - 0.02 | 0.03 | 0.16 - 0.60 |
| | 31 | 0.31 - 0.62 | 0.08 | 0.005 - 0.02 | 0.03 | 0.16 - 0.60 |
| | 32 | 0.32 - 0.64 | 0.08 | 0.005 - 0.02 | 0.03 | 0.17 - 0.64 |
| | 33 | 0.33 - 0.66 | 0.08 | 0.006 - 0.02 | 0.03 | 0.17 - 0.66 |
| | 34 | 0.34 - 0.68 | 0.09 | 0.006 - 0.02 | 0.03 | 0.17 - 0.68 |
| | 35 | 0.35 - 0.70 | 0.09 | 0.006 - 0.02 | 0.04 | 0.18 - 0.70 |

Rat Analgesic Dosages

| | Meloxicam | Meloxicam SR | Buprenorphine | Buprenorphine SR | Carprofen | Nocita |
|-----------------------------|----------------------------|--------------|---------------|------------------|----------------------------|--------------|
| Stock concentration (mg/ml) | 5 | 2 | 0.3 | 1 | 50 | 13.3 |
| instructions for diluting | 4 ml meloxicam + 6 ml NaCl | n/a | n/a | n/a | 1 ml carprofen + 9 ml NaCl | n/a |
| Diluted concentration for | 2 | n/a | n/a | n/a | 5 | n/a |
| dosage (mg/kg) | 2 | 4 | 0.1-0.5 | 1.0 - 1.2 | 5 | 5.3 |
| frequency | q 12 - 24 hr | q 72 hr | q 6 hr | q 72 hr | q 12 hr | once |
| RAT weight in grams | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) | Dose SQ (ml) |
| | 250 | 0.25 | 0.50 | 0.08 - 0.43 | 0.25 | 0.25 |
| | 260 | 0.26 | 0.52 | 0.09 - 0.45 | 0.26 | 0.26 |
| | 270 | 0.27 | 0.54 | 0.09 - 0.47 | 0.27 | 0.27 |
| | 280 | 0.28 | 0.56 | 0.09 - 0.48 | 0.28 | 0.28 |
| | 290 | 0.29 | 0.58 | 0.09 - 0.50 | 0.29 | 0.29 |
| | 300 | 0.30 | 0.60 | 0.1 - 0.52 | 0.30 | 0.30 |
| | 310 | 0.31 | 0.62 | 0.10 - 0.53 | 0.31 | 0.31 |
| | 320 | 0.32 | 0.64 | 0.11 - 0.55 | 0.32 | 0.32 |
| | 330 | 0.33 | 0.66 | 0.11 - 0.57 | 0.33 | 0.33 |
| | 340 | 0.34 | 0.68 | 0.11 - 0.58 | 0.34 | 0.34 |
| | 350 | 0.35 | 0.70 | 0.12 - 0.60 | 0.35 | 0.35 |
| | 360 | 0.36 | 0.72 | 0.12 - 0.62 | 0.36 | 0.36 |
| | 370 | 0.37 | 0.74 | 0.12 - 0.63 | 0.37 | 0.37 |
| | 380 | 0.38 | 0.76 | 0.13 - 0.65 | 0.38 | 0.38 |
| | 390 | 0.39 | 0.78 | 0.13 - 0.67 | 0.39 | 0.39 |
| | 400 | 0.40 | 0.80 | 0.13 - 0.68 | 0.40 | 0.40 |
| | 410 | 0.41 | 0.82 | 0.14 - 0.70 | 0.41 | 0.41 |
| | 420 | 0.42 | 0.84 | 0.14 - 0.72 | 0.42 | 0.42 |